Health Profile: *Cardiff, United Kingdom*

*Taking cities to a
healthier future*

EURO-URHIS 2

European Urban Health Indicators System Part 2
Urban Health Monitoring and Analysis System to Inform Policy

A	Skopje, the former Yugoslav Republic of Macedonia	N	Tromeø , Norway
B	Tetovo, the former Yugoslav Republic of Macedonia	O	Bistrița, Romania
C	Bordeaux, France	P	Craiova, Romania
D	Montpellier, France	Q	Iași, Romania
E	Oberhausen, Germany	R	Bratislava, Slovakia
F	Köln, Germany	S	Košice, Slovakia
G	Liepāja, Latvia	T	Ljubljana, Slovenia
H	Riga, Latvia	U	Maribor, Slovenia
I	Kaunas, Lithuania	V	Birmingham, United Kingdom
J	Šiauliai, Lithuania	W	Cardiff, United Kingdom
K	Amsterdam, The Netherlands	X	Glasgow, United Kingdom
L	Utrecht, The Netherlands	Y	Merseyside, United Kingdom
M	Oslo, Norway	Z	Greater Manchester, United Kingdom

Depression and anxiety were more often reported in Cardiff compared to the other EURO-URHIS 2 cities.

All-cause mortality in both males and females is similar in Cardiff compared to other EURO-URHIS 2 cities. Male mortality from malignant neoplasms and female mortality from diseases of the circulatory system are substantially lower than the overall EURO-URHIS 2 mean. Female mortality from malignant neoplasms and mortality from diseases of the respiratory system are substantially higher.

Heavy episodic drinking in Cardiff youth occurs as often as in other EURO-URHIS 2 cities, whereas binge drinking in adults occurs more often. Smoking in youth and adults occurs as often as in other EURO-URHIS 2 cities.

The proportion of overweight or obese adults is higher than the overall EURO-URHIS 2 proportion.

Health and health determinants in Cardiff vary considerably by age and level of education.

This health profile describes the health situation and associated health determinants in Cardiff compared with those observed in other European urban areas.

Cardiff is one of the urban areas chosen for EURO-URHIS 2 (European Urban Health Indicator System Part 2), a project that aims to identify health problems in urban areas. The EURO-URHIS 2 project describes health and health determinants specific to urban areas in Europe, covering cities in North, East, South, and West Europe. This project may add to information that is already locally available, in that it is the first study to enable reliable comparisons of health status between different cities in Europe. Policy makers can use the information to prioritise topics for urban health policy and for interventions in an evidence-based way.

EURO-URHIS 2 gathered information by collecting data from routinely available registration data, and by conducting youth and adult surveys at the end of 2010. In total, data from 26 urban areas in Europe were available for between-city comparisons and benchmarking.

The routinely available registration data relate to the most recently available year (2006-2008). The youth survey was a school-based survey of 14-16 year olds. In Cardiff, 304 students completed a valid questionnaire. The adult survey was carried out involving a representative sample of adults aged 19-64 and 65+. In Cardiff, 192 19-64 year olds and 252 65+ year olds completed valid questionnaires.

More detailed information on the justification of methods and instruments that were used, as well as response rates, selection of cities and indicators, and statistical methodology, can be found on our websites: www.urhis.eu and <http://results.urhis.eu>. The websites also provide data from other participating urban areas and comparisons between specific cities can be made.

The graphs in this health profile show the health status of the urban area compared to other EURO-URHIS 2 urban areas. The whiskers represent the lowest and highest value within the EURO-URHIS 2 project on a scale of 0 to 100%. The grey bar represents the 25th, 50th, and 75th percentile. The urban area value is shown as a diamond, which is blue when the value is not statistically significantly different from the EURO-URHIS 2 mean and red when the difference is statistically significant (at the 5% level).

Figure 1. Age distribution

Figure 2. Level of education

Differences in health status may possibly be explained by age and education. Figures 1 and 2 show the age distribution and level of education in Cardiff compared to the other EURO-URHIS 2 urban areas. Age and education did not explain any observed differences in the adult survey between Cardiff and other EURO-URHIS 2 urban areas.

DISCLAIMER

To achieve maximum quality of the data, all instruments used were based on knowledge of earlier studies and expert consultations, and were piloted, validated, and optimised. The survey questionnaires of EURO-URHIS 2 were based on already existing, validated instruments; selected indicators were as little culturally sensitive as possible. Questionnaires were translated in the local language(s) and, for validation purposes, back-translated into English. Youth survey response rates were generally very high. In the adult survey, a minimum response rate of 30% was required to be included for benchmarking. Despite all our efforts, and as in any survey, the point estimates for certain health indicators in your urban area may deviate from other estimates, and may not be comparable to other local information due to differences in study methodology and indicator definitions. If you would like further information regarding the methodology, please see our websites: <http://www.urhis.eu> and <http://results.urhis.eu>.

Health-related Characteristics of Cardiff

	Indicator	Cardiff	United Kingdom	EURO-URHIS 2 range (percentiles)					EURO-URHIS 2 mean	N
				min	25th	50th	75th	max		
Demographic	1. Population size (x1,000)	325	61,192	67	264	406	708	2,565	570	23
	2. Population density	2,176	251	27	1,115	2,040	2,840	4,580	1,974	24
	3. Population aged 0-19 years	24%	24%	17%	20%	22%	24%	28%	22%	23
	4. Population aged 65+ years	14%	16%	7%	11%	14%	15%	20%	14%	23
	5. Live births	58	63	39	45	52	58	75	53	24
	6. Teenage pregnancies	-	26	4	7	11	20	33	14	18
	7. Pregnancies after age 35	-	34	7	18	23	33	59	28	18
Socio-economic	8. Unemployment (age 19-64)	4.8%	-	3.6%	4.0%	4.9%	7.2%	10.2%	5.8%	16
	9. Higher level education	46%	-	25%	33%	45%	53%	72%	45%	16
	10. Not enough money	11%	-	5%	11%	16%	22%	61%	21%	16
	11. Low family wealth	14%	-	5%	7%	13%	21%	44%	16%	20
Health System	12. MMR vaccinated	88%	86%	83%	88%	94%	97%	100%	93%	19
	13. DTP vaccinated	96%	92%	83%	93%	95%	97%	99%	94%	19
	14. Cervical smear test	71%	-	41%	62%	70%	76%	83%	68%	16
	15. Cholesterol measurement	35%	-	23%	42%	47%	52%	64%	47%	16
Health Status	16. Life expectancy - male	76.6	77.8	68.2	71.0	75.3	76.1	77.0	73.6	18
	17. Life expectancy - female	81.7	81.9	76.2	78.5	80.2	81.0	82.0	79.7	18
	18. Infant mortality	2.6	4.6	1.3	3.5	4.9	5.7	9.4	5.0	24
	19. Low birth weight	7.3%	7.1%	2.7%	5.2%	6.6%	8.1%	11.8%	6.7%	22

Table 1. Health-related characteristics of Cardiff

Source. Indicators 1-7, 12-13, and 16-19: routinely available registration data; indicators 8-10 and 14-15: adult survey; indicator 11: youth survey. Missing data are indicated by "-".

N = number of urban areas that were able to collect data on the specific indicator.

1. number of inhabitants; **2.** number of inhabitants per km²; **3.** % of inhabitants aged 0-19 years; **4.** % of inhabitants aged 65 years or older; **5.** number of births per 1,000 women aged 15-44 years; **6.** number of births per 1,000 women aged 15-19 years; **7.** number of births per 1,000 women aged 35-44 years; **8.** % of adults aged 19-64 years who are unemployed; **9.** % of adults who attained higher level education; **10.** % of adults who do not have enough money for daily expenses; **11.** % of youth who live in a low wealth family, as defined by a FAS (Family Affluence Scale) score of ≤3; **12.** % of population who have completed measles, mumps, and rubella (MMR) vaccination courses before school-age; **13.** % of population who have completed diphtheria, tetanus, and poliomyelitis (DTP) vaccination courses before school-age; **14.** % of adult women who have undergone a cervical smear test within the past three years; **15.** % of adults who had their serum cholesterol measured within the last year; **16-17.** number of years that a newborn is expected to live if current mortality rates continue to apply; **18.** annual number of deaths of children under one year of age, per 1,000 births; **19.** % of total live births weighing less than 2,500 grams

Compared to other cities in EURO-URHIS 2, Cardiff is an urban area with average population density and a somewhat younger population.

The percentage of inhabitants with higher level education in Cardiff (46%) is similar to the overall EURO-URHIS 2 mean. The proportion of adults who reported to not have enough money for daily expenses (11%) and the proportion of youth that reported to live in poor families (14%) are similar to the other EURO-URHIS 2 urban areas.

The proportion of adults in Cardiff who had their serum cholesterol measured within the last year is significantly lower compared to the other EURO-URHIS 2 cities.

Life expectancy at birth is an indicator for the general health

status of a population. In Cardiff, male life expectancy is 76.6 years and female life expectancy is 81.7 years, which are both higher than the overall average in EURO-URHIS 2.

Infant mortality is an indicator for population health and quality of health care services. With an infant mortality rate of 2.6 per 1,000 live births, Cardiff belongs to the 25% of urban areas in EURO-URHIS 2 with the lowest infant mortality.

At the population level, low birth weight is an indicator for pregnancy conditions and perinatal care. Low birth weight can at the individual level also result in health problems later in life. Of all newborns in Cardiff, 7.3% had a low birth weight, which is comparable to the overall EURO-URHIS 2 mean.

YOUTH HEALTH STATUS

	Indicator	Cardiff	EURO-URHIS 2 range (percentiles)			EURO-URHIS 2 mean	N
			0%	50%	100%		
Health Status	1. Good self-perceived health	83%				92%	20
	2. Elevated risk of psychological problems	30%				20%	20
	3. Psychosomatic symptoms	14%				10%	20
	4. Low back pain	36%				42%	20
Lifestyle Factors	5. Overweight and obesity	-				13%	15
	6. Physical activity ≥2 hours/week	36%				50%	20
	7. Regular fruit consumption	37%				49%	20
	8. Regular vegetable/salad consumption	39%				52%	20
	9. Regular tooth brushing	76%				72%	20
	10. Frequently watching television	70%				60%	20
	11. Daily smoking	11%				12%	20
	12. First smoking ≤13 years	24%				24%	20
	13. Heavy episodic drinking	35%				33%	20
	14. First alcohol ≤13 years	52%				53%	19
	15. Ever used cannabis	21%				16%	20
	16. Unprotected sexual intercourse	7%				4%	20
Environ- ment	17. Crime in area	53%				35%	20
	18. Involved in traffic accident	6%				7%	18
	19. Being bullied	10%				7%	20

Table 2. Health status and determinants in youth (14-16 years)

Source. Indicators 1-19: youth survey. Missing data are indicated by "-".

N = number of urban areas that were able to collect data on the specific indicator.

1. % of youth who perceive their health as good, very good, or excellent; **2.** % of youth with an overall Strengths and Difficulties Questionnaire (SDQ) score of 20 or higher; **3.** % of youth who reported a lot of headaches, stomach aches, or sickness during the past six months; **4.** % of youth who experienced low back pain during the past month; **5.** % of youth overweight or obese according to the international BMI cut-offs; **6.** % of youth who participate in vigorous physical activity for more than two hours per week in their free time; **7.** % of youth who eat fruit on most days of the week; **8.** % of youth who eat vegetables and/or salads on most days of the week; **9.** % of youth who brush their teeth more than once a day; **10.** % of youth who watch television for more than two hours on weekdays; **11.** % of youth who smoke tobacco every day; **12.** % of youth who reported first smoking at ≤13 years; **13.** % of youth who drank five or more units of alcohol on one occasion during the past 30 days; **14.** % of youth who reported first drinking alcohol at ≤13 years; **15.** % of youth who ever used cannabis; **16.** % of the total youth population who did not use a condom the last time they had sexual intercourse; **17.** % of youth who reported presence of crime, violence, or vandalism in the area where they live; **18.** % of youth who had a road traffic accident resulting in injury over the past 12 months; **19.** % of youth who have been bullied at least twice in the past couple of months

Health Status and Determinants in Youth

Table 2 gives an overview of the health status and determinants in Cardiff youth, as reported from the survey. Self-perceived health is a measure of adolescent well-being. 83% of youth in Cardiff perceived their health to be (very) good or excellent, which is significantly lower than the overall EURO-URHIS 2 proportion. In Cardiff, a significantly higher proportion of youth were identified with an elevated risk of psychological problems (30%), compared to the overall EURO-URHIS 2 proportion. Psychosomatic symptoms like headaches, stomach aches, and sickness were reported more often in Cardiff, whereas low back pain was reported less often.

Childhood obesity is related to a higher risk of obesity, disability, and premature death later in life. Physical activity can contribute

to maintaining a healthy weight and preventing the occurrence of chronic conditions. Furthermore, physical activity is associated with psychological benefits and with a better school performance in young people. The proportion of youth who reported participation in vigorous physical activity for two or more hours per week is significantly lower in Cardiff (36%), compared to the overall EURO-URHIS 2 proportion. Sedentary behaviour is related to overweight and obesity, independent of physical activity. Youth in Cardiff watch significantly more television on weekdays compared to other urban areas in EURO-URHIS 2. A healthy diet can lower the risk of obesity. Regular consumption of fruit and vegetables occurs less frequently in Cardiff than in other EURO-URHIS 2 urban areas.

Initiation of smoking and drinking alcohol at a young age is a strong predictor of smoking during adulthood and of later problems with alcohol. The proportion of youth in Cardiff who smoke daily (11%) is similar to the overall EURO-URHIS 2 proportion. Heavy episodic drinking of five or more units of alcohol on one occasion was reported as often in Cardiff (35%) as in the total EURO-URHIS 2 population.

Regular cannabis use in young people can lead to impaired cognitive development. 21% of youth in Cardiff have ever used cannabis, which is higher than the overall EURO-URHIS 2 proportion.

The proportion of youth who reported they did not use a condom when they last had sexual intercourse is significantly higher in Cardiff compared to other EURO-URHIS 2 urban areas.

Neighbourhood crime, violence, or vandalism was significantly more often reported by youth in Cardiff (53%) compared to other cities. The proportion of youth who were victims of bullying in the past couple of months was significantly higher compared to the other urban areas in EURO-URHIS 2.

ADULT HEALTH STATUS

Indicator		Cardiff	United Kingdom	EURO-URHIS 2 range (percentiles)					EURO-URHIS 2 mean	N
				min	25th	50th	75th	max		
Morbidity	1. HIV/AIDS incidence - male	-	16*	2	6	8	23	71	16	19
	2. HIV/AIDS incidence - female	-	9*	0	2	6	12	16	7	19
	3. Tuberculosis incidence	15	14	5	11	17	39	153	33	22
	4. Lung cancer incidence	-	66	29	42	55	62	103	54	13
Mortality	5. All-cause mortality - male	755	729	654	752	834	1,014	1,426	919	19
	6. All-cause mortality - female	510	510	362	495	542	640	821	560	19
	7. Malignant neoplasms - male	230	216	195	230	245	258	336	250	22
	8. Malignant neoplasms - female	164	154	114	143	153	162	232	154	22
	9. Diseases of the circulatory system - male	249	247	154	227	298	456	676	353	22
	10. Diseases of the circulatory system - female	146	156	91	147	199	299	406	220	22
	11. Diseases of the respiratory system - male	87	89	32	55	62	80	158	72	22
	12. Diseases of the respiratory system - female	61	64	12	21	36	50	120	43	22
	13. Transport accidents	3	6	1	3	5	11	16	7	21
	14. Suicide and intentional harm	8	7	4	8	11	15	29	12	22

Table 3. Morbidity and mortality

Source. Indicators 1-14: routinely available registration data. Missing data are indicated by “-”.

* Country level data include HIV incidence only.

N = number of urban areas that were able to collect data on the specific indicator.

1-4. Number of newly diagnosed cases with a specific disease per 100,000 persons per year; **5-6.** All-cause mortality rate per 100,000 persons per year (standardised on European population); **7-14.** Mortality rate due to a specific cause per 100,000 persons per year (standardised on European population)

Health Status and Determinants in Adults

The health status of a population can be assessed by using a number of parameters, such as those referring to acute and chronic disease, mortality, psychological well-being, and self-perceived health. Table 3 and indicators 1-8 of Table 4 show the overall health status among adults in Cardiff, compared to other cities in Europe. The results show that in Cardiff the incidence of tuberculosis is similar to the overall average in all EURO-URHIS 2 urban areas.

Both for males and females, all-cause mortality is comparable to other cities. Mortality from malignant neoplasms is lower for males and higher for females. Mortality from diseases of the respiratory system is substantially higher, whereas female mortality from diseases of the circulatory system and mortality from suicide and intentional harm is lower than in other cities.

Health Status and Determinants in Adults (continued)

	Indicator	Cardiff	EURO-URHIS 2 range (percentiles)			EURO-URHIS 2 mean	N
			0%	50%	100%		
Health Status	1. (Very) good self-perceived health	67%				64%	16
	2. Psychological problems	24%				23%	16
	3. Depression/anxiety	16%				9%	16
	4. Cardiovascular disease (age 65+)	11%				18%	16
	5. Cancer	1%				2%	16
	6. Asthma or bronchitis	9%				7%	16
	7. Long-standing illness with restrictions	26%				28%	16
	8. Low back pain	34%				45%	16
Lifestyle Factors	9. Regular consumption of fruit/vegetables	64%				53%	16
	10. Regular breakfast	81%				78%	16
	11. Being physically active ≥ twice a week	42%				46%	16
	12. Overweight and obesity	56%				50%	16
	13. Daily smoking	15%				18%	16
	14. Passive smoking by non-smokers	7%				13%	16
	15. Binge drinking	25%				17%	16
	16. Cannabis last year (age 19-64)	10%				5%	16
Environment	17. Green areas suitable for recreational activities	90%				84%	16
	18. Belonging to immediate neighbourhood	53%				54%	16
	19. Social cohesion in neighbourhood	57%				52%	16
	20. Exposure to severe noise	8%				14%	16
	21. Damp spots or mould at home	34%				27%	16

Table 4. Health status and determinants in adults (19 years and older)

Source. Indicators 1-21: adult survey. Missing data are indicated by “-”.

N = number of urban areas that were able to collect data on the specific indicator.

1. % of adults who perceive their health to be good or very good; **2.** % of adults with a score of four or more on the General Health Questionnaire (GHQ); **3.** % of adults who reported to be diagnosed with or treated for anxiety or depression during the past year; **4.** % of adults aged 65 years and older who were diagnosed with or treated for heart attack, angina, or heart failure during the past year; **5.** % of adults who were diagnosed with or treated for (any kind of malignant) cancer during the past year; **6.** % of adults who were diagnosed with or treated for bronchial asthma or chronic bronchitis during the past year; **7.** % of adults who suffer from any long-standing illness, long-standing effect from injury, disability, or other long-standing condition; **8.** % of adults who had low back pain longer than one day in the past month; **9.** % of adults who eat, on average, four or more portions of fruit and/or vegetables per day; **10.** % of adults who have breakfast at least four times a week; **11.** % of adults who are physically active for at least 30 minutes twice a week or more; **12.** % of adults overweight or obese, defined as a BMI of ≥ 25 kg/m²; **13.** % of adults who smoke every day; **14.** % of non-smokers who are exposed to second-hand smoking inside their home; **15.** % of adults who drink six or more portions of alcohol on one occasion, at least once a week (men) or at least once a month (women); **16.** % of adults aged 19-64 years who used cannabis during the past year; **17.** % of adults who perceive the green areas in their neighbourhood to be suitable for active recreational activities; **18.** % of adults who feel that they belong to their immediate neighbourhood; **19.** % of adults who perceive their neighbourhood to be socially cohesive; **20.** % of adults who were exposed to severe noise from outdoors during the past 12 months; **21.** % of adults who had wet or damp spots and/or mould or mildew inside their homes (other than in basements) within the past 12 months

The proportion of people in Cardiff who perceive their health to be good or very good (67%) does not differ from the average in the other urban areas in EURO-URHIS 2. The percentage of adults who reported psychological problems in Cardiff (24%) is comparable to other urban areas in EURO-URHIS 2. Cancer, cardiovascular disease in the elderly, and low back pain were significantly less often reported in Cardiff. Depression and anxiety, on the contrary, are more prevalent.

Several lifestyle factors and environmental determinants can affect health (Table 4, indicators 9-21). Daily smoking, for instance, increases the risk of cancer, particularly lung cancer. Smokers are also at far greater risk of developing heart disease, stroke, and emphysema. Binge drinking is associated with many health problems, which include injuries and violence, sexually transmitted diseases, alcohol dependency, liver disease, and neurological damage. The percentage of persons who smoke

daily (15%) does not differ from other EURO-URHIS 2 cities, whereas exposure to second-hand smoking inside their home was less often reported in Cardiff. The proportion of adults who regularly drink more than six units of alcohol (25%) is significantly higher in Cardiff compared to the overall EURO-URHIS 2 mean. A significantly higher proportion of people in Cardiff reported to have used cannabis during the last year.

Being overweight and obese are important determinants of death worldwide. They increase the risk of cardiovascular diseases, diabetes, musculoskeletal disorders, and some cancers. In Cardiff, 56% of the adults are overweight or obese, which is higher than the overall EURO-URHIS 2 proportion. Being overweight and obese are related to lack of regular physical activity. Being physically active reduces the risk of hypertension, coronary heart disease, stroke, diabetes, breast and colon cancer, depression, and the risk of injury caused by falls.

The proportion of adults in Cardiff physically active more than twice a week (42%) does not differ from the total EURO-URHIS 2 proportion. A healthy diet can lower the risk of obesity. Adults in Cardiff more frequently eat fruit and vegetables.

Psychological well-being may be influenced both by the availability of green spaces in the neighbourhood that are suitable for recreational activities and by aspects of social cohesion. In Cardiff, 90% perceived their green spaces to be

suitable for recreational activities, whereas this proportion is significantly lower in other cities. The percentage of adults who perceived their neighbourhood to be socially cohesive was 57%, which is similar to the overall EURO-URHIS 2 average. A significantly lower percentage reported exposure to severe noise from outdoors, whereas the presence of damp spots or mould and mildew inside their homes was more often reported in Cardiff.

Indicator			Age		Gender		Education level	
		Total Population	19 - 64	65 +	Male	Female	Secondary level or lower	Higher level
Health Status	1. (Very) good self-perceived health	67%	69%*	57%*	62%	71%	57%*	78%*
	2. Psychological problems	24%	25%	20%	20%	28%	30%*	16%*
	3. Long-standing illness with restrictions	26%	23%*	41%*	24%	28%	36%*	15%*
Lifestyle Factors	4. Overweight and obesity	56%	55%	61%	61%	51%	59%	52%
	5. Daily smoking	15%	17%*	6%*	16%	14%	25%*	4%*
	6. Binge drinking	25%	28%*	10%*	29%	21%	23%	27%
	7. Regular consumption of fruit/vegetables	64%	62%*	74%*	59%	69%	58%	70%
	8. Being physically active ≥twice a week	42%	42%	38%	45%	39%	34%*	50%*
	9. Social cohesion in neighbourhood	57%	56%	64%	55%	60%	45%*	70%*

Table 5. Health and health determinants by demographic groups in Cardiff

Source. Adult survey.

Indicators are defined in Table 4. Missing data are indicated by "-".

* Statistically significant difference between subgroups at the 5% level.

Health and Health Determinants by Demographic Groups

Health and health determinants can vary considerably as according to age, gender, and education. Table 5 subdivides a selection of important health indicators in Cardiff by subgroup: respondents aged 19-64 and 65+ years, males and females, and adults who achieved secondary level education or lower and higher level education.

Respondents aged 19-64 years in Cardiff more often perceived their health to be good or very good, than is the case for older respondents. Younger respondents were less often restricted by a long-standing illness, but were more likely to be daily smokers or binge drinkers, and less frequently ate fruit and vegetables. The occurrence of psychological problems, overweight or obesity, physical activity, and perceived social neighbourhood cohesion did not differ by age.

Men and women in Cardiff did not differ in any of the studied indicators.

Adults in Cardiff who attained secondary level education or lower less often perceived their health to be good or very good and more frequently experienced psychological problems than adults with higher level education. Lower educated respondents were more often restricted by a long-standing illness, were more likely to be daily smokers, less often engaged in physical activity, and less often perceived their neighbourhood as being socially cohesive. The proportion of overweight or obesity, binge drinking, and fruit and vegetable consumption did not differ by education level.

Healthy Life Expectancy

Figure 3. Healthy Life Expectancy

Presented here are estimates of healthy life expectancy (HLE) at ages 15 and 60 for men and women in eligible EURO-URHIS 2 urban areas. HLE was calculated first by estimating life expectancy at each age using recent 5-year averages of all-causes mortality for each urban area. From this, life expectancy was broken down into years living in good and poor perceived health, estimated using responses to the EURO-URHIS 2 adult survey question: How is your health in general?: Very good/Good/Fair/Bad/Very bad/Don't know, and the youth survey question: In general, would you say your health is..?: Excellent/Very Good/Good/Fair/Poor. Those answering very good, good or fair on the adult survey were classed as being in good perceived health, with the remainder in poor perceived health. For the youth survey, fair and poor were categorised as poor perceived health to match the scale applied to the adult survey. It was then possible to calculate the total years in good and poor perceived health and present this as a population level HLE. Full details on this process will be available in the final EURO-URHIS 2 project report, available at www.urhis.eu.

Male life expectancy in Cardiff at age 15 was 60.1 years. This was 1.7 years less than the highest in the sample (Oslo, 61.8 years), and 7.3 years more than the lowest (Kaunas, 52.8 years). At this age, males were estimated to spend 55.1 years in good perceived health. This is 3.6 years less than the longest HLE (Amsterdam, 58.7 years) and 10.0 years more than the shortest (Siauliai, 45.1 years).

Male life expectancy in Cardiff at age 60 was 18.5 years. This was 1.1 years less than the highest in the sample (Oslo, 19.6 years), and 2.9 years more than the lowest (Siauliai, 15.6 years). At this age, males were estimated to spend 16.6 years in good perceived health. This is 1.1 years less than the longest HLE (Oslo, 17.7 years) and 5.8 years more than the shortest (Siauliai, 10.8 years).

Female life expectancy in Cardiff at age 15 was 62.8 years. This was 1.3 years less than the highest in the sample (Bordeaux, 64.1 years), and 1.4 years more than the lowest (Siauliai, 61.4 years). At this age, females were estimated to spend 55.5 years in good perceived health. This is 4.2 years less than the longest HLE (Oslo, 59.7 years) and 4.6 years more than the shortest (Kaunas, 50.9 years).

Female life expectancy in Cardiff at age 60 was 19.8 years. This was 1.1 years less than the highest in the sample (Bordeaux, 20.9 years), and 0.5 years more than the lowest (Siauliai, 19.3 years). At this age, females were estimated to spend 17.0 years in good perceived health. This is 2.3 years less than the longest HLE (Montpellier, 19.3 years) and 4.9 years more than the shortest (Siauliai, 12.1 years).

Landeszentrum Gesundheit
Nordrhein-Westfalen

Beneficiaries

The University of Manchester; Municipal Health Service Utrecht; University of Liverpool; The Iuliu Hatieganu University of Medicine & Pharmacy Epidemiology Department; The Norwegian Institute of Public Health; Municipal Health Service Amsterdam; Kaunas University of Medicine; Regional Public Health and Health Promotion Centre (Slovenia); Institute of Health and Work, North Rhine-Westphalia; Slovak Public Health Association; Hacettepe University, Department of Public Health; North West Regional Health Brussels Office; Latvian Public Health Agency; South East European University; National Federation of Regional Health Observatories; Pham Ngoc Thach University of Medicine

Local EURO-URHIS 2 representative in Cardiff:

Dr Arpana Verma
Manchester Academic Health Sciences Centre
University of Manchester

arpana.verma@manchester.ac.uk
+44 (0)161 275 5206

EURO-URHIS 2

European Urban Health Indicators System Part 2

Urban Health Monitoring and Analysis System to Inform Policy

The research leading to these results has received funding from the European Union's Seventh Framework Programme (FP7/2007-2013) under grant agreement no 223711

© EURO-URHIS 2, August 2012

Municipal Health Service Utrecht, The Netherlands

Design by Nicola Dale